

Request for Prior Authorization BIOLOGICALS FOR HIDRADENITIS SUPPURATIVA

FAX Completed Form To 1 (800) 574-2515 Provider Help Desk 1 (877) 776-1567

(PLEASE PRINT - ACCURACY IS IMPORTANT)

IA Medicaid Member ID #	Patient name		DOB
Patient address	,		
Provider NPI	Prescriber name		Phone
Prescriber address			Fax
Pharmacy name	Address		Phone
<u>-</u>	ormation above. It must be legible, corre	•	form will be returned.
Pharmacy NPI	Pharmacy fax	NDC	
Hidradenitis Suppurativa (HS which there is documentation Payment will be considered usefuested drug and indication interactions, and use in spectostage II or III disease; and 3) documentation of adequate to clindamycin; b) Oral clindamy for coverage are met, initial results.	quired for biologicals FDA approved.). Payment for non-preferred biologicals of a previous trial and therapy failurander the following conditions: 1) Ren, including age, dosing, contraindicial populations. 2) Patient has a diag Patient has at least three (3) abscess rials and therapy failures with the folycin plus rifampin; c) Maintenance the equests will be given for 4 months. A sal response to therapy. Clinical resp	c agents will be core with a preferred quest adheres to a ations, warnings a nosis of moderate ses or inflammator lowing: a) Daily tractable with a preference authorizational authorizational authorizational	onsidered only for cases in a biologic agent. all FDA approved labeling for and precautions, drug to severe HS with Hurley ry nodules; and 4) Patient has eatment with topical erred tetracycline. If criteria eations will be considered
total abscess and inflammato fistula count from initiation o	ory nodule count with no increase in f therapy. The required trials may be ents would be medically contraindic	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation o provided that use of these ag	ory nodule count with no increase in f therapy. The required trials may be	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation o provided that use of these ag	ory nodule count with no increase in f therapy. The required trials may be ents would be medically contraindic	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation o provided that use of these ag <u>Preferred</u>	ory nodule count with no increase in f therapy. The required trials may be lents would be medically contraindic lon-Preferred Bimzelx Cosentyx	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation of provided that use of these agon management of the search of the s	ory nodule count with no increase in f therapy. The required trials may be ents would be medically contraindic lon-Preferred Bimzelx	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation of provided that use of these ages of the	ory nodule count with no increase in f therapy. The required trials may be lents would be medically contraindic lon-Preferred Bimzelx Cosentyx	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation of provided that use of these agon in the second secon	ory nodule count with no increase in f therapy. The required trials may be lents would be medically contraindic lon-Preferred Bimzelx Cosentyx	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation of provided that use of these agon series and inflammator of these agon series and initiation of provided that use of these agon series and initiation of these agon series and initiation of the series and inflammator	ory nodule count with no increase in f therapy. The required trials may be lents would be medically contraindic lon-Preferred Bimzelx Cosentyx	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation of provided that use of these ages of the ages of these ages of these ages of the ages o	ory nodule count with no increase in f therapy. The required trials may be lents would be medically contraindic lon-Preferred Bimzelx Cosentyx	abscess count an overridden when	d no increase in draining
total abscess and inflammator fistula count from initiation of provided that use of these ages of the ages of these ages of the ages of these ages of the ag	ory nodule count with no increase in f therapy. The required trials may be tents would be medically contraindic lon-Preferred Bimzelx Cosentyx Other Humira Biosimilar:	abscess count an overridden when ated.	d no increase in draining documented evidence is
total abscess and inflammator fistula count from initiation of provided that use of these ages of the ages of these ages of these ages of these ages of the ages of these ages of the ages of th	ory nodule count with no increase in f therapy. The required trials may be tents would be medically contraindic lon-Preferred Bimzelx Cosentyx Other Humira Biosimilar:	abscess count an overridden when ated.	d no increase in draining documented evidence is
total abscess and inflammator fistula count from initiation of provided that use of these ages of the ages of these ages of the ages of these ages of these ages of these ages of the ages o	ory nodule count with no increase in f therapy. The required trials may be lents would be medically contraindic lon-Preferred Bimzelx Cosentyx Other Humira Biosimilar: Dosage Instructions	abscess count an overridden when ated. Quantity	d no increase in draining documented evidence is
total abscess and inflammator fistula count from initiation of provided that use of these ages Preferred	pry nodule count with no increase in f therapy. The required trials may be lents would be medically contraindic lon-Preferred Bimzelx Cosentyx Other Humira Biosimilar: Dosage Instructions	abscess count an overridden when ated. Quantity	d no increase in draining documented evidence is

470-5408 (Rev. 4/25) Page 1 of 2

Request for Prior Authorization BIOLOGICALS FOR HIDRADENITIS SUPPURATIVA

(PLEASE PRINT – ACCURACY IS IMPORTANT)

Topical Clindamycin Trial Name/Dose:			
Oral Clindamycin Plus Rifampin Trial:			
Clindamycin: Dose:	Trial dates:		
Reason for failure:			
Rifampin: Dose:	Trial dates:		
Reason for failure:			
Maintenance Preferred Tetracycline Trial:			
Name/Dose:	Trial dates:		
Reason for failure:			
Renewals			
Document response to therapy:			
Abscess/Nodule Count: Increase Decrease (provide count): Date obtained:			
Has patient had an increase in draining fistula count since initiation of the	erapy?		
Has patient had an increase in draining fistula count since initiation of the Other medical conditions to consider:			
·			
Other medical conditions to consider:			
Other medical conditions to consider: Possible drug interactions/conflicting drug therapies:			
Other medical conditions to consider:			

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary by contact with the county Department of Health and Human Services, that the member continues to be eligible for Medicaid.

470-5408 (Rev. 4/25) Page 2 of 2